

Dirac Equation in Gauge and Affine-Metric Gravitation Theories

GIOVANNI GIACHETTA

Department of Mathematics and Physics
University of Camerino, 62032 Camerino, Italy
E-mail: mangiarotti@camvax.unicam.it

GENNADI SARDANASHVILY

Department of Theoretical Physics
Moscow State University, 117234 Moscow, Russia
E-mail: sard@grav.phys.msu.su

Abstract

We show that the covariant derivative of Dirac fermion fields in the presence of a general linear connection on a world manifold is universal for Einstein's, gauge and affine-metric gravitation theories.

1

We follow the gauge approach to the description of gravitational interactions.

In reality, one observes three types of fields: gravitational fields, fermion fields and gauge fields associated with internal symmetries. We are not concerned here with Higgs fields whose dynamics remains elusive.

If the gauge invariance under internal symmetries is kept in the presence of a gravitational field, Lagrangian densities of gauge fields must depend on a metric gravitational field only.

Different spinor models of fermion matter have been suggested. At present, all observable fermion particles are Dirac fermions. Therefore, we restrict our consideration to Dirac fermion fields in gravitation theory.

In the gauge gravitation theory, gravity is represented by pairs (h, A_h) of gravitational fields h and associated Lorentz connections A_h [5, 10]. The connection A_h is usually identified with both a connection on a world manifold X and a spinor connection on the the spinor bundle $S_h \rightarrow X$ whose sections describe Dirac fermion fields ψ_h in the presence of the gravitational field h . The problem arises when Dirac fermion fields are described in the framework of the affine-metric gravitation theory. In this case, the fact that a world connection is some Lorentz connection may result from the field equations, but it can not be assumed in advance. There are models where the world connection is not a Lorentz connection [5]. Moreover, it may happen that a world connection is the Lorentz connection with respect to different gravitational fields [15]. At the same time, a Dirac fermion field can be regarded only in a pair (h, ψ_h) with a certain gravitational field h .

Indeed, one must define the representation of cotangent vectors to X by the Dirac's γ -matrices in order to construct the Dirac operator. Given a tetrad gravitational field $h(x)$, we have the representation

$$\gamma_h : dx^\mu \mapsto \hat{dx}^\mu = h_a^\mu \gamma^a.$$

However, different gravitational fields h and h' yield the nonequivalent representations γ_h and $\gamma_{h'}$.

It follows that, fermion-gravitation pairs (h, ψ_h) are described by sections of the composite spinor bundle

$$S \rightarrow \Sigma \rightarrow X \quad (1)$$

where $\Sigma \rightarrow X$ is the bundle of gravitational fields h where values of h play the role of parameter coordinates, besides the familiar world coordinates. [10, 13]. In particular, every spinor bundle $S_h \rightarrow X$ is isomorphic to the restriction of $S \rightarrow \Sigma$ to $h(X) \subset \Sigma$. Performing this restriction, we come to the familiar case of a field model in the presence of a gravitational field $h(x)$. The feature of the dynamics of field systems on the composite bundle (1) lies in the fact that we have the modified covariant differential of fermion fields which depend on derivatives of gravitational fields h .

As a consequence, we get the following covariant derivative of Dirac fermion fields in the presence of a gravitational field $h(x)$:

$$\widetilde{D}_\lambda = \partial_\lambda - \frac{1}{2} A^{abc}_\mu (\partial_\lambda h_c^\mu + K^\mu_{\nu\lambda} h_c^\nu) I_{ab}, \quad (2)$$

$$A^{abc}_\mu = \frac{1}{2} (\eta^{ca} h_\mu^b - \eta^{cb} h_\mu^a),$$

where K is a general linear connection on a world manifold X , η is the Minkowski metric, and

$$I_{ab} = \frac{1}{4} [\gamma_a, \gamma_b]$$

are generators of the spinor group $L_s = SL(2, \mathbf{C})$.

Let us emphasize that the connection

$$\widetilde{K}^{ab}_\lambda = A^{abc}_\mu (\partial_\lambda h_c^\mu + K^\mu_{\nu\lambda} h_c^\nu) \quad (3)$$

is not the connection

$$K^k_{m\lambda} = h_\mu^k (\partial_\lambda h_m^\mu + K^\mu_{\nu\lambda} h_m^\nu) = K^{ab}_\lambda (\eta_{am} \delta_b^k - \eta_{bm} \delta_a^k)$$

written with respect to the reference frame $h^a = h_\lambda^a dx^\lambda$, but there is the relation

$$\widetilde{K}^{ab}_\lambda = \frac{1}{2} (K^{ab}_\lambda - K^{ba}_\lambda). \quad (4)$$

If K is a Lorentz connection A_h , then the connection $\widetilde{K}$ (3) consists with K itself.

The covariant derivative (2) has been considered by several authors [1, 8, 16]. The relation (4) correspond to the canonical decomposition of the Lie algebra of the general linear group. By the well-known theorem [6], every general linear connection being projected onto the Lie algebra of the Lorentz group yields a Lorentz connection.

In our opinion, the advantage of the covariant derivative (2), consists in the fact that, being derived in the framework of the gauge gravitation theory, it may be also applied to the affine-metric gravitation theory and the conventional Einstein's gravitation theory. We are not concerned here with the general problem of equivalence of metric, affine and affine-metric theories of gravity [3, 7]. At the same time, when K is the Levi-Civita connection of h , the Lagrangian density of fermion fields which utilizes the covariant derivative (2) comes to that in the Einstein's gravitation theory. It follows that the configuration space of metric (or tetrad) gravitational fields and general linear connections may play the role of the universal configuration space of realistic gravitational models. In particular, one then can think of the generalized Komar superpotential as being the universal superpotential of energy-momentum of gravity [4].

2

We follow the geometric approach to field theory when classical fields are described by global sections of a bundle $Y \rightarrow X$ over a world manifold X . Their dynamics is phrased in terms of jet manifolds [11, 12, 14].

As a shorthand, one can say that the k -order jet manifold $J^k Y$ of a bundle $Y \rightarrow X$ comprises the equivalence classes $j_x^k s$, $x \in X$, of sections s of Y identified by the first $k+1$ terms of their Taylor series at a point x . Recall that a k -order differential operator on sections of a bundle $Y \rightarrow X$ is defined to be a bundle morphism of the bundle $J^k Y \rightarrow X$ to a vector bundle over X .

In particular, given bundle coordinates (x^μ, y^i) of a bundle Y , the first order jet manifold $J^1 Y$ of Y is endowed with the coordinates (x^μ, y^i, y_μ^i) where

$$y_\mu^i(j_x^1 s) = \partial_\mu s^i(x).$$

There is the 1:1 correspondence between the connections on the bundle $Y \rightarrow X$ and the global sections

$$\Gamma = dx^\lambda \otimes (\partial_\lambda + \Gamma_\lambda^i \partial_i)$$

of the affine jet bundle $J^1 Y \rightarrow Y$. Every connection Γ on $Y \rightarrow X$ yields the first order differential operator

$$\begin{aligned} D_\Gamma : J^1 Y &\xrightarrow{Y} T^* X \otimes_Y VY, \\ D_\Gamma &= (y_\lambda^i - \Gamma_\lambda^i) dx^\lambda \otimes \partial_i, \end{aligned}$$

on Y which is called the covariant differential relative to the connection Γ . We denote by VY the vertical tangent bundle of Y .

In the first order Lagrangian formalism, the first order jet manifold J^1Y of Y plays the role of the finite-dimensional configuration space of fields represented by sections s of a bundle $Y \rightarrow X$. A first order Lagrangian density is defined to be an exterior horizontal density

$$\begin{aligned} L : J^1Y &\rightarrow \bigwedge^n T^*X, & n &= \dim X, \\ L &= \mathcal{L}(x^\mu, y^i, y_\mu^i) \omega, & \omega &= dx^1 \wedge \dots \wedge dx^n, \end{aligned}$$

on the bundle $J^1Y \rightarrow X$.

Note that, since the jet bundle $J^1Y \rightarrow Y$ is affine, every polynomial Lagrangian density of field theory factors through

$$L : J^1Y \xrightarrow{D} T^*X \otimes_Y VY \rightarrow \bigwedge^n T^*X$$

where D is the covariant differential relative to some connection on Y .

3

Let us consider the gauge theory of gravity and fermion fields. By X is further meant an oriented 4-dimensional world manifold which satisfies the well-known topological conditions in order that gravitational fields and spinor structure can exist on X . To summarize these conditions, we assume that X is not compact and that the tangent bundle of X is trivial.

Let LX be the principal bundle of oriented linear frames in tangent spaces to X . In gravitation theory, its structure group $GL^+(4, \mathbf{R})$ is reduced to the connected Lorentz group $L = SO(1, 3)$. It means that there exists a reduced subbundle L^hX of LX whose structure group is L . In accordance with the well-known theorem, there is the 1:1 correspondence between the reduced L subbundles L^hX of LX and the global sections h of the quotient bundle

$$\Sigma := LX/L \rightarrow X. \tag{5}$$

These sections h describe gravitational fields on X , for the bundle (5) is the 2-fold covering of the bundle of pseudo-Riemannian metrics on X .

Given a section h of Σ , let Ψ^h be an atlas of LX such that the corresponding local sections z_ξ^h of LX take their values into L^hX . With respect to Ψ^h and a holonomic atlas $\Psi^T = \{\psi_\xi^T\}$ of LX , a gravitational field h can be represented by a family of GL_4 -valued tetrad functions

$$h_\xi = \psi_\xi^T \circ z_\xi^h, \quad dx^\lambda = h_a^\lambda(x) h^a. \tag{6}$$

By the Lorentz connections A_h associated with a gravitational field h are meant the principal connections on the reduced subbundle L^hX of LX . They give rise to principal connections on LX and to spinor connections on the L_s -lift P_h of L^hX .

There are different ways to introduce Dirac fermion fields. We follow the algebraic approach.

Given a Minkowski space M , let $Cl_{1,3}$ be the complex Clifford algebra generated by elements of M . A spinor space V is defined to be a minimal left ideal of $Cl_{1,3}$ on which this algebra acts on the left. We have the representation

$$\gamma : M \otimes V \rightarrow V$$

of elements of the Minkowski space $M \subset Cl_{1,3}$ by Dirac's matrices γ on V .

Let us consider a bundle of complex Clifford algebras $Cl_{1,3}$ over X whose structure group is the Clifford group of invertible elements of $Cl_{1,3}$. Its subbundles are both a spinor bundle $S_M \rightarrow X$ and the bundle $Y_M \rightarrow X$ of Minkowski spaces of generating elements of $Cl_{1,3}$. To describe Dirac fermion fields on a world manifold X , one must require Y_M to be isomorphic to the cotangent bundle T^*X of X . It takes place if there exists a reduced L subbundle $L^h X$ such that

$$Y_M = (L^h X \times M)/L.$$

Then, the spinor bundle

$$S_M = S_h = (P_h \times V)/L_s \quad (7)$$

is associated with the L_s -lift P_h of $L^h X$. In this case, there exists the representation

$$\gamma_h : T^*X \otimes S_h = (P_h \times (M \otimes V))/L_s \rightarrow (P_h \times \gamma(M \otimes V))/L_s = S_h \quad (8)$$

of cotangent vectors to a world manifold X by Dirac's γ -matrices on elements of the spinor bundle S_h . As a shorthand, one can write

$$\hat{d}x^\lambda = \gamma_h(dx^\lambda) = h_a^\lambda(x)\gamma^a.$$

Given the representation (8), we shall say that sections of the spinor bundle S_h describe Dirac fermion fields in the presence of the gravitational field h . Indeed, let

$$A_h = dx^\lambda \otimes (\partial_\lambda + \frac{1}{2}A^{ab}{}_\lambda I_{ab}{}^A{}_B \psi^B \partial_A)$$

be a principal connection on S_h . Given the corresponding covariant differential D and the representation γ_h (8), one can construct the Dirac operator

$$\mathcal{D}_h = \gamma_h \circ D : J^1 S_h \rightarrow T^*X \otimes_{S_h} V S_h \rightarrow V S_h, \quad (9)$$

$$\dot{y}^A \circ \mathcal{D}_h = h_a^\lambda \gamma^{aA}{}_B (y_\lambda^B - \frac{1}{2}A^{ab}{}_\lambda I_{ab}{}^A{}_B y^B)$$

on the spinor bundle S_h .

Different gravitational fields h and h' define nonequivalent representations γ_h and $\gamma_{h'}$. It follows that a Dirac fermion field must be regarded only in a pair with a certain gravitational field. There is the 1:1 correspondence between these pairs and sections of the composite spinor bundle (1).

4

By a composite bundle is meant the composition

$$Y \rightarrow \Sigma \rightarrow X. \quad (10)$$

of a bundle $Y \rightarrow X$ denoted by Y_Σ and a bundle $\Sigma \rightarrow X$. It is coordinatized by $(x^\lambda, \sigma^m, y^i)$ where (x^μ, σ^m) are coordinates of Σ and y^i are the fiber coordinates of Y_Σ . We further assume that Σ has a global section.

The application of composite bundles to field theory is founded on the following [9]. Given a global section h of Σ , the restriction Y_h of Y_Σ to $h(X)$ is a subbundle of $Y \rightarrow X$. There is the 1:1 correspondence between the global sections s_h of Y_h and the global sections of the composite bundle (10) which cover h . Therefore, one can think of sections s_h of Y_h as describing fermion fields in the presence of a background parameter field h , whereas sections of the composite bundle Y describe all the pairs (s_h, h) . The configuration space of these pairs is the first order jet manifold $J^1 Y$ of the composite bundle Y .

The feature of the dynamics of field systems on composite bundles consists in the following. Every connection

$$A_\Sigma = dx^\lambda \otimes (\partial_\lambda + \tilde{A}_\lambda^i \partial_i) + d\sigma^m \otimes (\partial_m + A_m^i \partial_i)$$

on the bundle Y_Σ yields the horizontal splitting

$$VY = VY_\Sigma \oplus_Y (Y \times V\Sigma),$$

$$\dot{y}^i \partial_i + \dot{\sigma}^m \partial_m = (\dot{y}^i - A_m^i \dot{\sigma}^m) \partial_i + \dot{\sigma}^m (\partial_m + A_m^i \partial_i).$$

Building on this splitting, one can construct the first order differential operator

$$\begin{aligned} \widetilde{D} : J^1 Y &\rightarrow T^* X \otimes_Y VY_\Sigma, \\ \widetilde{D} &= dx^\lambda \otimes (y_\lambda^i - \tilde{A}_\lambda^i - A_m^i \sigma_\lambda^m) \partial_i, \end{aligned} \quad (11)$$

on the composite bundle Y . This operator possesses the following property.

Given a global section h of Σ , let Γ be a connection on Σ whose integral section is h , that is, $\Gamma \circ h = J^1 h$. It is readily observed that the differential (11) restricted to $J^1 Y_h \subset J^1 Y$ comes to the familiar covariant differential relative to the connection

$$A_h = dx^\lambda \otimes [\partial_\lambda + (A_m^i \partial_\lambda h^m + \tilde{A}_\lambda^i) \partial_i]$$

on Y_h . Thus, it is $\widetilde{D}$ that we may utilize in order to construct a Lagrangian density

$$L : J^1 Y \xrightarrow{\widetilde{D}} T^* X \otimes_Y VY_\Sigma \rightarrow \wedge^n T^* X$$

for sections of the composite bundle Y .

5

In gravitation theory, we have the composite bundle

$$LX \rightarrow \Sigma \rightarrow X$$

where Σ is the quotient bundle (5) and

$$LX_\Sigma := LX \rightarrow \Sigma$$

is the L -principal bundle.

Let P_Σ be the L_s -principal lift of LX_Σ such that

$$P_\Sigma/L_s = \Sigma, \quad LX_\Sigma = r(P_\Sigma).$$

In particular, there is the imbedding of the L_s -lift P_h of $L^h X$ onto the restriction of P_Σ to $h(X)$.

Let us consider the composite spinor bundle (1) where

$$S_\Sigma = (P_\Sigma \times V)/L_s$$

is associated with the L_s -principal bundle P_Σ . It is readily observed that, given a global section h of Σ , the restriction S_Σ to $h(X)$ is the spinor bundle S_h (7) whose sections describe Dirac fermion fields in the presence of the gravitational field h .

Let us provide the principal bundle LX with a holonomic atlas $\{\psi_\xi^T, U_\xi\}$ and the principal bundles P_Σ and LX_Σ with associated atlases $\{z_\epsilon^s, U_\epsilon\}$ and $\{z_\epsilon = r \circ z_\epsilon^s\}$. With respect to these atlases, the composite spinor bundle is endowed with the bundle coordinates $(x^\lambda, \sigma_a^\mu, \psi^A)$ where $(x^\lambda, \sigma_a^\mu)$ are coordinates of the bundle Σ such that σ_a^μ are the matrix components of the group element $(\psi_\xi^T \circ z_\epsilon)(\sigma)$, $\sigma \in U_\epsilon$, $\pi_{\Sigma X}(\sigma) \in U_\xi$. Given a section h of Σ , we have

$$(\sigma_a^\lambda \circ h)(x) = h_a^\lambda(x),$$

where $h_a^\lambda(x)$ are the tetrad functions (6).

Let us consider the bundle of Minkowski spaces

$$(LX \times M)/L \rightarrow \Sigma$$

associated with the L -principal bundle LX_Σ . Since LX_Σ is trivial, it is isomorphic to the pullback $\Sigma \times_X T^*X$ which we denote by the same symbol T^*X . Then, one can define the bundle morphism

$$\gamma_\Sigma : T^*X \otimes_{\Sigma} S_\Sigma = (P_\Sigma \times (M \otimes V))/L_s \rightarrow (P_\Sigma \times \gamma(M \otimes V))/L_s = S_\Sigma, \quad (12)$$

$$\hat{d}x^\lambda = \gamma_\Sigma(dx^\lambda) = \sigma_a^\lambda \gamma^a,$$

over Σ . When restricted to $h(X) \subset \Sigma$, the morphism (12) comes to the morphism γ_h (8).

We use this morphism in order to construct the total Dirac operator on the composite spinor bundle S (1).

Let

$$\tilde{A} = dx^\lambda \otimes (\partial_\lambda + \tilde{A}_\lambda^B \partial_B) + d\sigma_a^\mu \otimes (\partial_\mu^a + A_\mu^{Ba} \partial_B)$$

be a principal connection on the bundle S_Σ and $\tilde{D}$ the corresponding differential (11). We have the first order differential operator

$$\mathcal{D} = \gamma_\Sigma \circ \tilde{D} : J^1 S \rightarrow T^* X \otimes_S V S_\Sigma \rightarrow V S_\Sigma,$$

$$\dot{\psi}^A \circ \mathcal{D} = \sigma_a^\lambda \gamma^{aA}_B (\psi_\lambda^B - \tilde{A}_\lambda^B - A_\mu^{Ba} \sigma_{a\lambda}^\mu),$$

on S . One can think of it as being the total Dirac operator since, for every section h , the restriction of $\mathcal{D}$ to $J^1 S_h \subset J^1 S$ comes to the Dirac operator $\mathcal{D}_h$ (9) relative to the connection

$$A_h = dx^\lambda \otimes [\partial_\lambda + (\tilde{A}_\lambda^B + A_\mu^{Ba} \partial_\lambda h_a^\mu) \partial_B]$$

on the bundle S_h .

In order to construct the differential $\tilde{D}$ (11) on $J^1 S$ in explicit form, let us consider the principal connection on the bundle LX_Σ which is given by the local connection form

$$\tilde{A} = (\tilde{A}^{ab}_\mu dx^\mu + A^{abc}_\mu d\sigma_c^\mu) \otimes I_{ab}, \quad (13)$$

$$\tilde{A}^{ab}_\mu = \frac{1}{2} K^\nu_{\lambda\mu} \sigma_c^\lambda (\eta^{ca} \sigma_\nu^b - \eta^{cb} \sigma_\nu^a),$$

$$A^{abc}_\mu = \frac{1}{2} (\eta^{ca} \sigma_\mu^b - \eta^{cb} \sigma_\mu^a), \quad (14)$$

where K is a general linear connection on TX and (14) corresponds to the canonical left-invariant free-curvature connection on the bundle

$$GL^+(4, \mathbf{R}) \rightarrow GL^+(4, \mathbf{R})/L.$$

Accordingly, the differential $\tilde{D}$ relative to the connection (13) reads

$$\tilde{D} = dx^\lambda \otimes [\partial_\lambda - \frac{1}{2} A^{abc}_\mu (\sigma_{c\lambda}^\mu + K^\mu_{\nu\lambda} \sigma_c^\nu) I_{ab}^A \psi^B \partial_A]. \quad (15)$$

Given a section h , the connection $\tilde{A}$ (13) is reduced to the Lorentz connection $\tilde{K}$ (3) on $L^h X$, and the differential (15) leads to the covariant derivatives of fermion fields (2).

We utilize the differential (15) in order to construct a Lagrangian density of Dirac fermion fields. Their Lagrangian density is defined on the configuration space $J^1(S \oplus_\Sigma S^+)$ coordinatized by

$$(x^\mu, \sigma_a^\mu, \psi^A, \psi_A^+, \sigma_{a\lambda}^\mu, \psi_\lambda^A, \psi_{A\lambda}^+).$$

It reads

$$\begin{aligned}
L_\psi = & \left\{ \frac{i}{2} [\psi_A^+ (\gamma^0 \gamma^\lambda)^A_B (\psi_\lambda^B - \frac{1}{2} A^{abc}_\mu (\sigma_{c\lambda}^\mu + K^\mu_{\nu\lambda} \sigma_c^\nu) I_{ab}{}^B{}_C \psi^C) - \right. \\
& \left. (\psi_{A\lambda}^+ - \frac{1}{2} A^{abc}_\mu (\sigma_{c\lambda}^\mu + K^\mu_{\nu\lambda} \sigma_c^\nu) \psi_C^+ I_{ab}{}^{+C}{}_A) (\gamma^0 \gamma^\lambda)^A_B \psi^B] - \right. \\
& \left. m \psi_A^+ (\gamma^0)^A_B \psi^B \right\} \sigma^{-1} \omega
\end{aligned} \tag{16}$$

where

$$\gamma^\mu = \sigma_a^\mu \gamma^a, \quad \sigma = \det(\sigma_a^\mu),$$

and

$$\psi_A^+ (\gamma^0)^A_B \psi^B$$

is the Lorentz invariant fiber metric in the bundle $S \oplus_\Sigma S^*$ [2].

One can easily verify that

$$\frac{\partial \mathcal{L}_\psi}{\partial K^\mu_{\nu\lambda}} + \frac{\partial \mathcal{L}_\psi}{\partial K^\mu_{\lambda\nu}} = 0.$$

Hence, the Lagrangian density (16) depends on the torsion of the general linear connection K only. In particular, it follows that, if K is the Levi-Civita connection of a gravitational field $h(x)$, after the substitution $\sigma_c^\nu = h_c^\nu(x)$, the Lagrangian density (16) comes to the familiar Lagrangian density of fermion fields in the Einstein's gravitation theory.

Acknowledgement

The authors would like to thank Prof. L.Mangiarotti and Prof. Y.Obukhov for valuable discussions.

References

1. A.Aringazin and A.Mikhailov, Matter fields in spacetime with vector non-metricity, *Clas. Quant. Grav.* **8**, 1685 (1991).
2. J.Crawford, Clifford algebra: Notes on the spinor metric and Lorentz, Poincaré and conformal groups, *J. Math. Phys.* **32**, 576 (1991).
3. M.Ferraris and J.Kijowski, On the equivalence of the relativistic theories of gravitation, *GRG* **14**, 165 (1982).
4. G.Giachetta and G.Sardanashvily, Stress-energy-momentum of affine-metric gravity. Generalized Komar superpotential, E-print: gr-qc/9511008.
5. F.Hehl, J.McCrea, E.Mielke and Y.Ne'eman: Metric-affine gauge theory of gravity, *Phys. Rep.* **258**, 1 (1995).

6. S.Kobayashi and K.Nomizu, *Foundations of Differential Geometry, Vol.1* (John Wiley, N.Y. - Singapore, 1963).
7. G.Magnano, Are there metric theories of gravity other than General Relativity, E-print: gr-qc/9511027.
8. V.Ponomarev and Yu.Obukhov, Generalized Einstein-Maxwell theory, *GRG* **14**, 309 (1982).
9. G.Sardanashvily, On the geometry of spontaneous symmetry breaking, *J. Math. Phys.* **33**, 1546 (1992).
10. G.Sardanashvily and O. Zakharov, *Gauge Gravitation Theory*, (World Scientific, Singapore, 1992).
11. G.Sardanashvily, *Gauge Theory in Jet Manifolds* (Hadronic Press, Palm Harbor, 1993).
12. G.Sardanashvily, Five lectures on the jet manifold methods in field theory, E-print: hep-th/9411089.
13. G.Sardanashvily, Composite spinor bundles in gravitation theory, E-print: gr-qc/9502022.
14. D.Saunders, *The Geometry of Jet Bundles* (Cambridge Univ. Press, Cambridge, 1989).
15. G.Thompson, Non-uniqueness of metrics compatible with a symmetric connection, *Class. Quant. Grav.*, **10**, 2035 (1993).
16. R.Tucker and C.Wang, Black holes with Weyl charge and non-Riemannian waves, *Class. Quant. Grav.* **12**, 2587 (1995).